

Duality and KPZ in Liouville Quantum Gravity

Bertrand Duplantier¹ and Scott Sheffield²

¹*Institut de Physique Théorique, CEA/Saclay, F-91191 Gif-sur-Yvette Cedex, France*

²*Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139**

(Dated: January 2, 2009)

We present a (mathematically rigorous) probabilistic and geometrical proof of the KPZ relation between scaling exponents in a Euclidean planar domain D and in Liouville quantum gravity. It uses the properly regularized quantum area measure $d\mu_\gamma = \varepsilon^{\gamma^2/2} e^{\gamma h_\varepsilon(z)} dz$, where dz is Lebesgue measure on D , γ is a real parameter, $0 \leq \gamma < 2$, and $h_\varepsilon(z)$ denotes the mean value on the circle of radius ε centered at z of an instance h of the Gaussian free field on D . The proof extends to the boundary geometry. The singular case $\gamma > 2$ is shown to be related to the quantum measure $d\mu_{\gamma'}$, $\gamma' < 2$, by the fundamental duality $\gamma\gamma' = 4$.

PACS numbers: 02.50.-r, 02.90.+p, 04.60.-m, 04.60.Kz, 05.40.-a, 11.25.Hf, 64.60.al

Introduction. One of the major theoretical advances in physics over the past thirty years has been the realization in gauge theory or string theory that transition amplitudes require summing over random surfaces, which replaces traditional sums over random paths, i.e., the celebrated Feynman path integrals of quantum mechanics and quantum field theory. In a 1981 breakthrough, A.M. Polyakov [1, 2] first understood that the summation over random Riemannian metrics involved could be represented mathematically by the so-called *Liouville theory of quantum gravity*.

The latter can be simply described as follows: Consider a bounded planar domain $D \subset \mathbb{C}$ as the parameter domain of the random Riemannian surface, and an instance h of the Gaussian free field (GFF) on D , with Dirichlet energy $(h, h)_\nabla := (2\pi)^{-1} \int_D \nabla h(z) \cdot \nabla h(z) dz$. The quantum area is then (formally) defined by $\mathcal{A} = \int_D e^{\gamma h(z)} dz$, where dz is the standard Euclidean (i.e., Lebesgue) measure and $e^{\gamma h(z)}$ the (random) conformal factor of the Riemannian metric, with a constant $0 \leq \gamma < 2$. The quantum Liouville action is then

$$S(h) = \frac{1}{2}(h, h)_\nabla + \lambda \mathcal{A}, \quad (1)$$

where $\lambda \geq 0$ is the so-called “cosmological constant”.

In 1986, V.A. Kazakov introduced the idea of placing a (critical) statistical model on a random planar lattice, when exactly solving there the Ising model [3]. This led to the 1988 breakthrough by Knizhnik, Polyakov and Zamolodchikov [4], building on [5], who predicted that corresponding critical exponents (i.e., conformal weights x) in the Euclidean plane and in quantum gravity (Δ) would obey the KPZ relation:

$$x = \frac{\gamma^2}{4} \Delta^2 + \left(1 - \frac{\gamma^2}{4}\right) \Delta. \quad (2)$$

In the critical continuum limit, the statistical system born by the random lattice is described by a conformal field theory (CFT) with central charge $c \leq 1$, which fixes

$\gamma = (\sqrt{25-c} - \sqrt{1-c})/\sqrt{6} \leq 2$ [4]. In Liouville quantum gravity the CFT action is then simply added to the Liouville one (1), allowing an alternative derivation of (2) in the so-called conformal gauge [6, 7].

This provides the core continuous model of “two-dimensional quantum gravity”, whose connections to string theory and conformal field theory, and to random planar lattice models and random matrix theory are manifold; see, e.g., the surveys [2, 8, 9, 10, 11, 12, 13] and references therein.

Despite the fact that the original work by KPZ has been cited almost a thousand times, and relation (2) repeatedly checked, in particular by explicit calculations in geometrical models on random planar lattices [14, 15, 16, 17], the KPZ relation was not proven rigorously until very recently. In the probabilistic and geometrical approach of [18], it is proven by starting from the *critical* Liouville gravity, with action S (1) taken at $\lambda = 0$, i.e., a *free-field* action, and (2) appears as a consequence of the fine properties of the two-dimensional GFF. Several follow-up works appeared since, either at the rigorous level [19, 20] or heuristic one [21].

One striking and important consequence of our perspective is that KPZ holds not only within the original CFT realm, but for *any fractal structure as measured with the random metric or stochastic measure $e^{\gamma h(z)} dz$, and for any $0 \leq \gamma < 2$* . For instance, it suggests that the same Euclidean exponent x of a random or self-avoiding walk would obey (2) with $\gamma = \sqrt{8/3}$ in pure gravity ($c = 0$), but with $\gamma = \sqrt{3}$ on a random lattice equilibrated with Ising spins ($c = 1/2$), yielding different Δ 's.

The aim of this Letter is to present this derivation in a minimal, yet rigorous way to the physics community. It involves using a properly defined and regularized quantum area measure, which allows for a transparent geometrical understanding of KPZ. It utilizes only the underlying Brownian stochastic properties of the GFF, whose correlation functions can be expressed in terms of the classical two-dimensional Newtonian potential. We also prove the boundary analog of KPZ for fractal sub-




FIG. 1: Potential f_ε^z (3) created by a uniform mass distribution on the circle $\partial B_\varepsilon(z)$; its leading constant value inside the disc $B_\varepsilon(z)$ is $-\log \varepsilon$.

sets of the boundary ∂D [18].

Our probabilistic approach also allows us to consider the *duality* property of Liouville quantum gravity: for $\gamma > 2$, the *singular* quantum measure can be properly defined in terms of the *regular* γ' -quantum measure, for the dual value $\gamma' = 4/\gamma < 2$, confirming that the so-called “other branch” of the γ -KPZ relation corresponds to standard γ' -KPZ for $\gamma' < 2$, as advocated long ago by Klebanov [22, 23, 24].

GFF circular average and Brownian motion. Let h be a centered Gaussian free field on a bounded simply connected domain D with Dirichlet zero boundary conditions. As already remarked in [5], special care is required to make sense of the quantum gravity measure, since the GFF is a distribution and not a function (it typically oscillates between $\pm\infty$) (see e.g., [25]).

For each $z \in D$, write $B_\varepsilon(z) = \{w : |w - z| < \varepsilon\}$. When $B_\varepsilon(z) \subset D$, write $h_\varepsilon(z)$ for the average value of h on the circle $\partial B_\varepsilon(z)$. Denote by $\rho_\varepsilon^z(y)$ the uniform Dirac density (of total mass one) on the circle $\partial B_\varepsilon(z)$, such that one can write the scalar product on D : $h_\varepsilon(z) = (h, \rho_\varepsilon^z) := \int_D h(y) \rho_\varepsilon^z(y) dy$. To the density ρ_ε^z is naturally associated a Newtonian potential. We define the function $f_\varepsilon^z(y)$, for $y \in D$,

$$f_\varepsilon^z(y) = -\log \max(\varepsilon, |z - y|) - \tilde{G}_z(y) \quad (3)$$

where $\tilde{G}_z(y)$ is the *harmonic* function of $y \in D$, with boundary value equal to the restriction of $-\log |z - y|$ to ∂D . By construction this $f_\varepsilon^z(y)$ satisfies Dirichlet b.c.’s and the Poisson equation $-\Delta f_\varepsilon^z = 2\pi \rho_\varepsilon^z$. This (regular) potential function is represented in Fig.1. Integrating by parts, we immediately have the following:

$$h_\varepsilon(z) = (h, f_\varepsilon^z)_\nabla, \quad (4)$$

in terms of the Dirichlet inner product

$$(f_1, f_2)_\nabla := (2\pi)^{-1} \int_D \nabla f_1(z) \cdot \nabla f_2(z) dz,$$

i.e., the interaction energy of fields associated with potentials $f_{i=1,2}$. In fact, the random variables $(h, f)_\nabla$ are zero mean Gaussian random variables for each f ,

with the covariance property: $\text{Cov}((h, f_1)_\nabla, (h, f_2)_\nabla) = (f_1, f_2)_\nabla$. From this, we deduce the covariance of the averaged $h_\varepsilon(z)$ fields (4) on two nested circles (Fig.1): $\text{Cov}(h_\varepsilon(z), h_{\varepsilon'}(z)) = (f_\varepsilon^z, f_{\varepsilon'}^z)_\nabla$. The latter is the Newtonian interaction energy of the two circles, which, owing to the explicit potential (3), to Gauss’ theorem and to harmonicity of $\tilde{G}_z$, gives the explicit form of the covariance

$$\mathbb{E}(h_\varepsilon(z) h_{\varepsilon'}(z)) = -\log \max(\varepsilon, \varepsilon') - \tilde{G}_z(z), \quad (5)$$

with $\mathbb{E} h_\varepsilon(z) = 0$ for Dirichlet boundary conditions, and where $\tilde{G}_z(z) = -\log C(z; D)$ in terms of the so-called conformal radius C of D viewed from z , a smooth function of z . From (5) we thus get the two important variances

$$\text{Var } h_\varepsilon(z) = -\log \varepsilon + \log C(z; D) \quad (6)$$

$$\text{Var}[h_\varepsilon(z) - h_{\varepsilon'}(z)] = |\log \varepsilon - \log \varepsilon'|. \quad (7)$$

The interpretation of (6,7) is immediate: for fixed z , the Gaussian random variable $h_\varepsilon(z)$ is *one-dimensional standard Brownian motion when parameterized by time* $t := -\log \varepsilon$ [18].

Random metrics and Liouville quantum gravity. Recall first that if N is a Gaussian random variable with mean a and variance b then $\mathbb{E} e^N = e^{a+b/2}$. Since $\mathbb{E} h_\varepsilon(z) = 0$, we have from (6) the exponential expectation:

$$\mathbb{E} e^{\gamma h_\varepsilon(z)} = e^{\text{Var}[\gamma h_\varepsilon(z)]/2} = [C(z; D)/\varepsilon]^{\gamma^2/2}. \quad (8)$$

Since (8) ultimately *diverges* for $\varepsilon \rightarrow 0$, we are led to *regularize* Liouville quantum gravity by defining the random measure

$$d\mu_\gamma = M_\varepsilon(z) dz, \quad M_\varepsilon(z) := \varepsilon^{\gamma^2/2} e^{\gamma h_\varepsilon(z)}, \quad (9)$$

in a way similar to the so-called Wick normal ordering (see e.g., [26]). In [18], it is shown that the limit of this regularized measure exists as $\varepsilon \rightarrow 0$, which mathematically defines Liouville quantum gravity (see also [27]).

GFF sampling and random metrics. We now consider a measure on pairs (z, h) where h is the Gaussian free field, and given h the point z is chosen from the regularized quantum area measure $e^{\gamma h_\varepsilon(z)} dz$. Such a measure has the form $e^{\gamma h_\varepsilon(z)} dh dz$ where dh represents the (whole) GFF measure. Its total action is thus the quadratic combination $\frac{1}{2}(h, h)_\nabla - \gamma h_\varepsilon(z)$. Owing to (4) and to $\text{Var } h_\varepsilon(z) = (f_\varepsilon^z, f_\varepsilon^z)_\nabla$, the latter can be rewritten as $\frac{1}{2}(h', h')_\nabla - \frac{\gamma^2}{2} \text{Var } h_\varepsilon(z)$, with the substitution $h' := h - \gamma f_\varepsilon^z$. The probability weight involved in our random metric can therefore be written as

$$\exp \left[-\frac{1}{2}(h', h')_\nabla \right] \mathbb{E} e^{\gamma h_\varepsilon(z)}, \quad (10)$$

where the second factor is the marginal distribution density (8) of z . The meaning of (10) is that, after sampling

z from its marginal distribution, the law of h weighted by $e^{\gamma h_\varepsilon(z)}$ is identical to that of the original GFF h' plus the deterministic function γf_ε^z (3).

KPZ proof. It is shown in [18] that when ε is small, the stochastic quantum measure (9) of the Euclidean ball $B_\varepsilon(z)$ is very well approximated by

$$\mu_\gamma(B_\varepsilon(z)) \simeq \pi \varepsilon^{\gamma Q} e^{\gamma h_\varepsilon(z)}, \quad (11)$$

where $Q_\gamma := 2/\gamma + \gamma/2$. In the simplified perspective of this work, we take (11) to be the *definition* of $\mu_\gamma(B_\varepsilon(z))$. That is, we view μ_γ as a function on balls of the form $B_\varepsilon(z)$, defined by (11), rather than a fully defined measure on D . Let us then call *quantum ball* $\tilde{B}^\delta(z)$ of area δ centered at z the (largest) Euclidean ball $B_\varepsilon(z)$ whose radius ε is chosen so that

$$\mu_\gamma(\tilde{B}^\delta(z)) = \delta. \quad (12)$$

One says that a (deterministic or random) fractal subset X of D has *Euclidean scaling exponent* x (and Euclidean dimension $2 - 2x$) if, for z chosen uniformly in D and independently of X , the probability $\mathbb{P}\{B_\varepsilon(z) \cap X \neq \emptyset\} \asymp \varepsilon^{2x}$, in the sense that $\lim_{\varepsilon \rightarrow 0} \log \mathbb{P} / \log \varepsilon = 2x$. Similarly, we say that X has *quantum scaling exponent* Δ if when X and (z, h) , sampled with weight (10), are chosen independently we have

$$\mathbb{P}\{\tilde{B}^\delta(z) \cap X \neq \emptyset\} \asymp \delta^\Delta. \quad (13)$$

In weight (10), $h' = h - \gamma f_\varepsilon^z$ is a standard GFF, thus its average has the characteristic property (7): $\mathcal{B}_t := h'_{\varepsilon=e^{-t}}(z)$ is standard Brownian motion in time $t = -\log \varepsilon$. Eq. (3) then gives $h_\varepsilon(z) = h'_\varepsilon(z) - \gamma \log \varepsilon = \mathcal{B}_t + \gamma t$ (up to a bounded constant), i.e., $h_\varepsilon(z)$ in (11) sampled with (10) has the same law as *Brownian motion with drift*.

Equality of (11) to (12) then relates stochastically the Euclidean radius ε to the quantum area δ . This radius is given in terms of the *stopping time*

$$T_A = -\log \varepsilon_A := \inf\{t : -\mathcal{B}_t + a_\gamma t = A\} \quad (14)$$

with the definitions $A := -(\log \delta)/\gamma > 0$ and $a_\gamma := Q_\gamma - \gamma = 2/\gamma - \gamma/2 > 0$ for $\gamma < 2$. A constant is absorbed in the choice of time origin such that $\mathcal{B}_0 = 0$.

The probability that the ball $B_{\varepsilon_A}(z)$ intersects X scales as $\varepsilon_A^{2x} = e^{-2xT_A}$. Computing its expectation $\mathbb{E} \exp(-2xT_A)$ with respect to the random time T_A will give the quantum probability (13). Consider then for any β the standard *Brownian exponential martingale* $\mathbb{E}[\exp(-\beta \mathcal{B}_t - \beta^2 t/2)] = 1$, valid for $0 \leq t < \infty$. We can apply it at the stopping time T_A , when $T_A < \infty$ and where $\mathcal{B}_{T_A} = a_\gamma T_A - A$; we thus get for $2x = \beta_\gamma a_\gamma + \beta_\gamma^2/2$

$$\begin{aligned} \mathbb{E}[\exp(-2xT_A) 1_{T_A < \infty}] &= \exp(-\beta_\gamma A) = \delta^{\Delta_\gamma} \\ \beta_\gamma(x) &:= (a_\gamma^2 + 4x)^{1/2} - a_\gamma, \quad \Delta_\gamma := \beta_\gamma/\gamma. \end{aligned} \quad (15)$$

For $x = 0$ one finds in particular $\mathbb{P}(T_A < \infty) = \mathbb{E}[1_{T_A < \infty}] = 1$, since $\beta_\gamma(0) = 0$ for $a_{\gamma < 2} > 0$, so that the conditioning on $T_A < \infty$ can actually be omitted. We thus obtain the expected quantum scaling behavior (13) with $\Delta = \Delta_\gamma$, which is the positive root to KPZ (2), **QED**.

The inverse Laplace transform $P_A(t)$ of (15), with respect to $2x$, is the probability density of $T_A = -\log \varepsilon_A$ such that $P_A(t)dt := \mathbb{P}(T_A \in [t, t+dt])$ [18]:

$$P_A(t) = \frac{A}{\sqrt{2\pi t^3}} \exp\left[-\frac{1}{2t}(A - a_\gamma t)^2\right]. \quad (16)$$

From (16) one deduces that for A large (i.e., δ and ε small), $\frac{A}{T_A} = \frac{\log \delta}{\gamma \log \varepsilon_A}$ is concentrated in (15) near $a_\gamma + \gamma \Delta$. Reverse engineering to GFF h via (14), one finds that a point z that is typical with respect to the quantum measure is an α -thick point of h [28]: $\alpha := \lim_{\varepsilon \rightarrow 0} \log h_\varepsilon(z) / \log \varepsilon^{-1} = \gamma - \gamma \Delta$, for a fractal of quantum scaling dimension Δ .

Boundary KPZ. Suppose that D is a domain with a (piecewise) linear boundary ∂D and h a GFF, now with *free* boundary conditions. For $z \in \partial D$, $h_\varepsilon(z)$ is the mean value of $h(z)$ on the *semicircle* $\partial B_\varepsilon(z) \cap D$, with variance scaling like $-2 \log \varepsilon$. We define the boundary quantum measure $d\mu_\gamma^B := \varepsilon^{\gamma^2/4} e^{\gamma h_\varepsilon(z)/2} dz$, where now dz is Lebesgue measure on ∂D , with the conformal factor needed for integrating a quantum length instead of an area, and a regulator such that the limit of μ_γ^B exists for $\varepsilon \rightarrow 0$ and $\gamma < 2$ [18]. For a fractal $X \subset \partial D$, we define boundary Euclidean ($\tilde{x}$) and quantum ($\tilde{\Delta}$) scaling exponents with this measure. We can repeat the analysis above, with now $h_\varepsilon(z)$ a standard Brownian motion $\mathcal{B}_{2t}$, with drift $\gamma t = -\gamma \log \varepsilon$, and prove the validity of KPZ relation (2) for the pair $(\tilde{x}, \tilde{\Delta})$ [18], as anticipated in [13].

Liouville quantum duality. For $\gamma > 2$, the Liouville measure (9) corresponds to the so-called “other” gravitational dressing of the Liouville potential [22, 23, 24]. The corresponding random surface is meant to be the scaling limit of random simply connected surfaces with large amounts of area cut off by small bottlenecks [29, 30, 31, 32, 33, 34, 35]. This surface turns out to be a tree-like foam of Liouville quantum bubbles of *dual* parameter $\gamma' := 4/\gamma$ (“baby-universes”) connected to each other at “pinch points” and rooted at a “principle bubble” parameterized by D . A precise description requires additional machinery and will appear elsewhere. For now we relate γ to γ' only formally.

The definition of quantum balls in (11) and (12) makes sense when $\gamma > 2$. Noting that $Q_{\gamma'} = Q_\gamma$, we have

$$\mu_{\gamma'}(B_\varepsilon(z)) = \pi \varepsilon^{\gamma' Q} e^{\gamma' h_\varepsilon(z)} = \mu_\gamma(B_\varepsilon(z))^{\gamma'/\gamma} = \mu_\gamma^{4/\gamma^2}$$

(up to an irrelevant power of π) — i.e., a γ -quantum ball of size δ has γ' -quantum size $\delta' := \delta^{4/\gamma^2}$. (Intuitively, the ball contains about a δ' fraction of the total γ' -quantum area but only a $\delta < \delta'$ fraction of the γ -quantum area

because the latter also includes points on non-principle bubbles.) The number of γ -quantum size- δ balls needed to cover the principle bubble D thus scales as $(\delta')^{-1} = \delta^{-4/\gamma^2}$.

From (2), the quantum scaling exponent Δ_γ in (15), when generalized to $\gamma > 2$, satisfies the *duality relation* $\Delta_\gamma - 1 = (4/\gamma^2)(\Delta_{\gamma'} - 1)$ [13, 22, 23, 24]. If $X \subset D$ has scaling exponent x , then (13), established for $\gamma' < 2$, essentially says (see [18]) that the expected number $N_{\gamma'}(\delta', X)$ of γ' -quantum size- δ' balls (*i.e.*, number $N_\gamma(\delta, X)$ of γ -quantum size- δ balls) required to cover X scales as $(\delta')^{\Delta_{\gamma'}-1} = \delta^{\Delta_\gamma-1}$.

Brownian approach to duality. When $\gamma > 2$, the ε -regularized measures $M_\varepsilon(z)dz$ (9) converge to zero. If we choose the pair (z, h) from the weighted measure $M_\varepsilon(z)dhdz$ as in (10) and consider the Brownian description (14), we find that $a_\gamma < 0$ for $\gamma > 2$, *i.e.*, the drift term runs in a direction opposite to $A > 0$, so that $T_A = \infty$ for large A . The weighted measure is thus *singular*; *i.e.*, there is a quantum area of at least δ localized at z for small enough δ . The Brownian martingale result (15) for $x = 0$ gives the probability, at a given z , for T_A to be finite:

$$\mathbb{P}(T_A < \infty) = \mathbb{E}[1_{T_A < \infty}] = \delta^{\Delta_\gamma(0)} = \delta^{1-4/\gamma^2} = \delta/\delta',$$

where $\Delta_\gamma(0) = (|a_\gamma| - a_\gamma)/\gamma = 1 - 4/\gamma^2$. For general x , (15) scales as δ^{Δ_γ} . We may define a δ -regularized measure $M^\delta(z)dhdz$ as $M_{\varepsilon_A}(z)dhdz$ restricted to the event $T_A < \infty$. Replacing γ with $\gamma' = 4/\gamma$ and δ with δ' has the same effect as multiplying $M^\delta(z)$ by $\delta'/\delta = \delta^{4/\gamma^2-1}$, so $\frac{\delta'}{\delta}M^\delta(z)dz$ converges to $d\mu_{\gamma'}$. This agrees with the conditional expectation scaling as

$$\frac{\mathbb{E}[\exp(-2xT_A)1_{T_A < \infty}]}{\mathbb{E}[1_{T_A < \infty}]} = \delta^{\Delta_\gamma} \times \frac{\delta'}{\delta} = \delta'^{\Delta_{\gamma'}}.$$

Using (15) and $a_{\gamma'} = -a_\gamma$ one obtains $\Delta_\gamma \Delta_{\gamma'} = x$, as anticipated in [13]. The typical GFF thickness $\alpha = \gamma(1 - \Delta_\gamma) = \gamma'(1 - \Delta_{\gamma'})$ is *invariant* under duality and obeys the Seiberg bound $\alpha \leq Q$ [8]; the *string susceptibility exponent* $\gamma_{\text{str}} = 2 - 2Q/\gamma$ obeys the expected duality relation $(1 - \gamma_{\text{str}})(1 - \gamma'_{\text{str}}) = 1$ [13, 22, 23, 24, 29, 31, 32, 33, 34, 35]. Finally, for the SLE $_\kappa$ process $\gamma = \sqrt{\kappa}$ [18], so that the Liouville $\gamma\gamma' = 4$ and SLE $\kappa\kappa' = 16$ dualities coincide.

We thank O. Angel, J. Franchi, P. Jones, I. Klebanov, I. Kostov, G.-F. Lawler, A. Okounkov and the late Oded Schramm for useful discussions.

- [1] A. M. Polyakov, Phys. Lett. B **103**, 207 (1981).
- [2] A. M. Polyakov, *Gauge fields and strings* (Harwood Academic Publishers, Chur, 1987).
- [3] V. A. Kazakov, Phys. Lett. A **119**, 140 (1986).
- [4] V. G. Knizhnik, A. M. Polyakov, and A. B. Zamolodchikov, Mod. Phys. Lett. A **3**, 819 (1988).
- [5] A. M. Polyakov, Mod. Phys. Lett. A **2**, 893 (1987).
- [6] F. David, Mod. Phys. Lett. A **3**, 1651 (1988).
- [7] J. Distler and H. Kawai, Nucl. Phys. B **321**, 509 (1989).
- [8] N. Seiberg, Progr. Theoret. Phys. Suppl. **102**, 319 (1990).
- [9] P. Ginsparg and G. Moore, in *Recent direction in particle theory, Proceedings of the 1992 TASI*, edited by J. Harvey and J. Polchinski (World Scientific, Singapore, 1993).
- [10] F. David, in *Gravitation et quantifications (Les Houches, Session LVII, 1992)*, edited by B. Julia and J. Zinn-Justin (Elsevier B.V., Amsterdam, 1995), pp. 679–749.
- [11] J. Ambjørn, B. Durhuus, and T. Jonsson, *Quantum geometry* (C.U.P., Cambridge, 1997).
- [12] Y. Nakayama, Int. J. Mod. Phys. A **19**, 2771 (2004).
- [13] B. Duplantier, in *Fractal geometry and applications: a jubilee of Benoît Mandelbrot, Part 2* (Amer. Math. Soc., Providence, RI, 2004), vol. 72 of *Proc. Sympos. Pure Math.*, pp. 365–482, arXiv:math-ph/0303034.
- [14] B. Duplantier and I. Kostov, Phys. Rev. Lett. **61**, 1433 (1988).
- [15] I. K. Kostov, Mod. Phys. Lett. A **4**, 217 (1989).
- [16] V. A. Kazakov and I. K. Kostov, Nucl. Phys. B **386**, 520 (1992).
- [17] B. Duplantier, Phys. Rev. Lett. **81**, 5489 (1998).
- [18] B. Duplantier and S. Sheffield (2008), arXiv:0808.1560.
- [19] I. Benjamini and O. Schramm (2008), arXiv:0806.1347.
- [20] R. Rhodes and V. Vargas (2008), arXiv:0807.1036.
- [21] F. David and M. Bauer (2008), arXiv:0810.2858.
- [22] I. R. Klebanov, Phys. Rev. D **51**, 1836 (1995).
- [23] I. R. Klebanov and A. Hashimoto, Nucl. Phys. B **434**, 264 (1995).
- [24] I. R. Klebanov and A. Hashimoto, Nucl. Phys. B Proc. Suppl. **45**, 135 (1996).
- [25] S. Sheffield, Probab. Th. Rel. Fields **139**, 521 (2007).
- [26] B. Simon, *The $P(\Phi)_2$ Euclidean (quantum) field theory* (P.U.P., Princeton, 1974).
- [27] R. Høegh-Krohn, Comm. Math. Phys. **21**, 244 (1971).
- [28] X. Hu, J. Miller, and Y. Peres, unpublished (2008).
- [29] S. R. Das, A. Dhar, A. M. Sengupta, and S. R. Wadia, Mod. Phys. Lett. A **5**, 1041 (1990).
- [30] S. Jain and S. D. Mathur, Phys. Lett. B **286**, 239 (1992).
- [31] G. P. Korchemsky, Mod. Phys. Lett. A **7**, 3081 (1992).
- [32] G. P. Korchemsky, Phys. Lett. B **296**, 323 (1992).
- [33] L. Alvarez-Gaumé, J. L. F. Barbón, and Č. Crnković, Nucl. Phys. B **394**, 383 (1993).
- [34] B. Durhuus, Nucl. Phys. B **426**, 203 (1994).
- [35] J. Ambjørn, B. Durhuus, and T. Jonsson, Mod. Phys. Lett. A **9**, 1221 (1994).

* On leave of absence from the Courant Institute for Mathematical Sciences at NYU.